

Exclusive proton asymmetry measurement in non-mesonic weak decay of polarized ${}^5_{\Lambda}\text{He}$.

T. Maruta,¹ S. Ajimura,² K. Aoki,³ A. Banu,⁴ H. Bhang,⁵ T. Fukuda,^{3,*} O. Hashimoto,⁶ J. I. Hwang,⁵ S. Kameoka,⁶ B. H. Kang,⁵ E. H. Kim,⁵ J. H. Kim,^{5,†} M. J. Kim,⁵ Y. Miura,⁶ Y. Miyake,² T. Nagae,³ M. Nakamura,¹ S. N. Nakamura,⁶ H. Noumi,³ S. Okada,^{7,‡} Y. Okayasu,⁶ H. Outa,^{3,‡} H. Park,⁸ P. K. Saha,^{3,§} Y. Sato,³ M. Sekimoto,³ T. Takahashi,^{6,¶} H. Tamura,⁶ K. Tanida,^{1,‡} A. Toyoda,³ K. Tsukuda,⁶ T. Watanabe,⁶ and H. J. Yim⁵

¹*Department of Physics, University of Tokyo, Tokyo 113-0033, Japan*

²*Department of Physics, Osaka University, Osaka 560-0043, Japan*

³*High Energy Accelerator Research Organization (KEK), Tsukuba, Ibaraki 305-0801, Japan*

⁴*GSI, Darmstadt D-64291, Germany*

⁵*Department of Physics, Seoul National University, Seoul 151-742, Korea*

⁶*Physics Department, Tohoku University, Sendai 980-8578, Japan*

⁷*Department of Physics, Tokyo Institute of Technology, Tokyo 152-8551, Japan*

⁸*Korea Research Institute of Standards and Science (KRISS), Daejeon, 305-600, Korea*

(Dated: February 8, 2008)

The asymmetry parameter α_p^{NM} for a proton exclusively emitted in the $\Lambda p \rightarrow np$ process was, for the first time, measured in the non-mesonic weak decay of a polarized ${}^5_{\Lambda}\text{He}$ hypernucleus by selecting the proton-neutron pairs emitted in the back-to-back kinematics. The highly polarized ${}^5_{\Lambda}\text{He}$ was abundantly produced with the (π^+, K^+) reaction at 1.05 GeV/c in the scattering angular range of $\pm 15^\circ$. The obtained value $\alpha_p^{NM} = 0.31 \pm 0.22$, as well as that for inclusive protons, $\alpha_p^{NM} = 0.11 \pm 0.08 \pm 0.04$, largely contradicts recent theoretical values of around -0.6 , although these calculations well reproduce the branching ratios of non-mesonic weak decay.

PACS numbers: 21.80.+a, 13.30.Eg, 13.75.Ev

In free space, a Λ hyperon mainly decays through a mesonic decay process ($\Lambda \rightarrow N\pi$), in which the momentum transfer (Δq) is about 100 MeV/c. On the other hand, the non-mesonic weak decay process (NMWD : $\Lambda N \rightarrow NN$) becomes dominant because of the large momentum transfer ($\Delta q \sim 400$ MeV/c). Since it is experimentally very difficult to carry out two-body scattering measurements in $\Lambda N \rightarrow NN$ or $NN \rightarrow \Lambda N$ reactions at this moment, studying the NMWDs of Λ hypernuclei gives us a very unique opportunity to investigate a new type of baryon-baryon weak interaction. It also has an advantage that both parity conserving and parity non-conserving parts can be studied through the flavor-changing weak process; the former part is usually masked by the strong interaction in the case of NN weak processes.

There have been two experimental observables used to investigate the NMWD process. One is the ratio of the partial decay width between neutron-induced decay, Γ_n ($\Lambda n \rightarrow nn$), and that of proton-induced decay, Γ_p ($\Lambda p \rightarrow np$), called as the Γ_n/Γ_p ratio. It is sensitive to the isospin structure of the ΛN weak interaction, because it allows only the final isospin, $I_f = 1$, in neutron-induced decay, while $I_f = 0$ and 1 are allowed in proton-induced decay. The other is the asymmetry parameter of the decay proton from the NMWD process (α_p^{NM}). It is obtained from the angular distribution $W(\theta)$ of decay particles emitted from polarized hypernuclei, as follows:

$$W(\theta) \propto 1 + A \cos \theta = 1 + \alpha P_{\Lambda} \cos \theta, \quad (1)$$

TABLE I: Six channels of NMWD start from the ΛN relative S-states.

ΛN	NN	amplitude	I_f	parity conservation
${}^1S_0$	${}^1S_0$	a^2	1	yes
	${}^3P_0$	b^2	1	no
${}^3S_1$	${}^3S_1$	c^2	0	yes
	${}^3D_1$	d^2	0	yes
	${}^1P_1$	e^2	0	no
	${}^3P_1$	f^2	1	no

where A is the asymmetry, P_{Λ} denotes the polarization of a Λ hypernucleus and θ is the emission angle of decay particles with respect to the polarization axis.

The asymmetry parameter comes from the interference between different final parity and isospin states. For example, concerning the s -shell hypernucleus ${}^5_{\Lambda}\text{He}$, the decay amplitudes are classified to six amplitudes according to the final spin states of the NN system, as listed in Table I [1]. Three amplitudes (a , b , f) have the final-state isospin $I_f = 1$, and the others (c , d , e) have $I_f = 0$. Then, the asymmetry parameter α_p^{NM} is expressed as [2]

$$\alpha_p^{NM} = \frac{2\sqrt{3}\text{Re}[ae^* - \frac{1}{\sqrt{3}}b(c - \sqrt{2}d)^* + f(\sqrt{2}c + d)^*]}{|a|^2 + |b|^2 + 3(|c|^2 + |d|^2 + |e|^2 + |f|^2)}. \quad (2)$$

So far, much attention has been paid to the former one Γ_n/Γ_p both experimentally and theoretically, because early measurements suggested Γ_n/Γ_p values $\gtrsim 1$,

while naive theoretical estimates based on a one-pion-exchange model gave small values of around 0.1 [3]. This large discrepancy seems to be resolved with a new measurement [4] and new theoretical calculations that take account of kaon exchange etc. [5, 6]; both now agree with the Γ_n/Γ_p to be $0.4 \sim 0.5$. However, it has been realized that such calculations happen to predict large negative values of around $-0.6 \sim -0.7$ for the decay asymmetry parameter α_p^{NM} . This significantly contradicts with a measured α_p^{NM} value of 0.24 ± 0.22 for ${}^5_\Lambda\text{He}$ by Ajimura *et al.* in the KEK E278 experiment [7].

There could be several effects to reduce the amplitude of the measured value. One of the effects is the final-state interaction (FSI) for nucleons emitted in the $\Lambda N \rightarrow NN$ processes. The detected proton could not be a proton directly emitted from the $\Lambda p \rightarrow np$ process, but secondary protons that underwent by FSIs, for which the initial asymmetry information would be lost. Another effect is the possible contamination of a two-nucleon induced decay process ($\Lambda p N \rightarrow np N$) in which the proton decay asymmetry would be very small because emitted nucleons are assumed to be distributed in three-body phase space uniformly. These protons from the secondary processes and two-nucleon induced process should have lower energies compared with the proton energy in the direct process. A previous measurement set a rather high detection threshold at ~ 60 MeV for protons to avoid possible contaminations of those low-energy protons. However, the ambiguity on the contaminations left a substantial systematic error.

In the present experiment KEK E462, we aimed at a high-statistics measurement of α_p^{NM} for ${}^5_\Lambda\text{He}$ together with a high detection efficiency for neutrons. It enabled us not only to measure the α_p^{NM} for protons directly coming from the $\Lambda p \rightarrow np$ process by requiring a neutron emitted in the opposite direction, but also to investigate the proton energy-threshold dependence of the α_p^{NM} values.

The experiment was performed at the K6 beam line of the KEK 12-GeV proton synchrotron (PS) with the high-resolution and large-acceptance superconducting kaon spectrometer (SKS) [8]. A polarized Λ hypernucleus ${}^5_\Lambda\text{He}$ was produced in the ${}^6\text{Li}(\pi^+, K^+)$ reaction at 1.05 GeV/c [9] in the scattering angle range of $\pm 15^\circ$ in the horizontal reaction plane. In the reaction, the ground state of ${}^6_\Lambda\text{Li}$, which is proton unbound, is initially produced, and soon decays into ${}^5_\Lambda\text{He}$ by emitting a proton. Therefore, the production of the ${}^5_\Lambda\text{He}$ was assured by tagging the production of the ground state of ${}^6_\Lambda\text{Li}$ in the (π^+, K^+) missing-mass spectrum.

The polarization of the ${}^5_\Lambda\text{He}$ hypernucleus was measured from the decay asymmetry of π^- s A_π emitted from the mesonic weak decay of ${}^5_\Lambda\text{He}$ by using two decay counter systems symmetrically installed above and below the Li target. Since the ${}^5_\Lambda\text{He}$ is composed of an α cluster and a Λ hyperon, it would be a good approxi-

mation to assume that the asymmetry parameter of the mesonic weak decay α_π^M is almost the same as that in free space [11], i.e. $\alpha_\pi^M = -0.642 \pm 0.013$. Then, the polarization of the ${}^5_\Lambda\text{He}$ is obtained from the measured decay asymmetry A_π .

The proton decay asymmetry A_p was measured at the same time by using the same decay counter systems.

Each decay counter system consisted of a set of drift chambers, two sets of timing counters and a neutron counter array with six layers of plastic counters. Both charged and neutral decay particles were identified by them. Concerning charged particles, pions, protons and deuterons, which were observed for the first time, were clearly separated. Neutrons were well-separated from gamma rays and we selected neutrons with the energy from 5 MeV to 150 MeV. The details of the experimental setup and particle identification are described in Ref. 10.

By selecting the ground-state region from -4 to 4 MeV in excitation energy, we succeeded to detect 5.2×10^4 events of ${}^5_\Lambda\text{He}$ in the ground state. The statistics are several-times higher than that of the previous experiment.

In fact, the asymmetry A in Eq. 1 was obtained from the up/down ratio between the yield in the up counter system N_U and that in the down counter system N_D . Owing to the large acceptance of the SKS spectrometer, we can polarize the hypernucleus upward (downward) by selecting the (π^+, K^+) scattering direction on the left (right) in one setup. Thus, we can take the double ratio R as

$$R = \left(\frac{N_U^+ \times N_D^-}{N_U^- \times N_D^+} \right)^{\frac{1}{2}} = \frac{1 + A \cos(0 + \varepsilon)}{1 + A \cos(\pi - \varepsilon)} = \frac{1 + A\eta}{1 - A\eta}, \quad (3)$$

where $N_{U(D)}^{+(-)}$ presents the yield in the up (down) counter system at the scattering angle to the left (right). Since the decay counter systems have some finite solid angles, the average measured angle $\langle \theta \rangle$ can not be 0 and π , and is shifted by some fraction ε . In this expression, systematic errors coming from the difference of the detection efficiencies and the acceptances between the up and down decay counter systems are canceled out. This first-order cancellation was examined in an analysis with the (π^+, pX) reaction, where X is either a proton or a pion, simultaneously obtained with the (π^+, K^+) reaction. No asymmetry is expected for protons and pions in this reaction and we confirmed it within an error of less than 0.3% for all the kaon scattering angles, as listed in Table II.

At first, the polarization of ${}^5_\Lambda\text{He}$ P_Λ was obtained as $P_\Lambda = A_\pi/\alpha_\pi^M$ from the mesonic decay process of the ground state, in which α_π^M is assumed to be almost the same as that in free space. The obtained polarizations P_Λ 's are listed in Table II. They are larger than the previous result [12], particularly in the large scattering angle region.

Then, using these P_Λ values, α_p^{NM} was obtained from




FIG. 1: Result of proton asymmetries for different scattering angles. Since the horizontal axis denotes the polarization of ${}^5_\Lambda\text{He}$ and the vertical one is the proton asymmetry, α_p^{NM} is obtained from a linear fit of two scattering angle regions, $6^\circ < |\theta_K| < 9^\circ$ and $9^\circ < |\theta_K| < 15^\circ$.

the asymmetry of proton A_p from the NMWD process of the ${}^5_\Lambda\text{He}$ ground state as $\alpha_p^{NM} = A_p/P_\Lambda$. Figure 1 shows the result of the asymmetries of decay proton as a function of the polarization P_Λ measured at three different kaon scattering angles in the Lab. system (θ_K): $2^\circ < |\theta_K| < 6^\circ$, $6^\circ < |\theta_K| < 9^\circ$ and $9^\circ < |\theta_K| < 15^\circ$. From a linear fitting, the α_p^{NM} value was obtained to be 0.07 ± 0.08 for $6^\circ < |\theta_K| < 15^\circ$. The systematic error was estimated by assuming the maximum pion contamination into protons. Since the pion asymmetry parameter has a large negative value ($\alpha^M = -0.642 \pm 0.013$), the contamination tends to reduce α_p^{NM} . Even by taking account of this effect, we obtained α_p^{NM} to be $0.11 \pm 0.07 \pm 0.04$, and confirmed that α_p^{NM} is slightly positive and small with better statistical accuracy.

Up to this point, all of the protons detected in the decay counter systems, of which detection energy threshold was about 30 MeV, were included in the analysis. As already discussed, there could be some protons not directly emitted from the $\Lambda p \rightarrow np$ process, which might reduce the observed asymmetry of protons. In Fig. 2 (a), the opening angle distribution between a proton and a neutron emitted from the ${}^5_\Lambda\text{He}$ in coincidence is shown. The peak structure at $\cos\theta_{np} \sim -1$ is attributed to the proton-neutron pairs directly emitted from the $\Lambda p \rightarrow np$ process, as expected from two-body kinematics. An excitation energy spectrum with proton-neutron pairs in the back-to-back direction ($\cos\theta_{np} < -0.8$) is also shown in Fig. 2 (b). The ground-state peak was clearly identified with very small background. The proton decay asymmetry parameter α_p^{NM} was obtained with these back-to-back coincidence events with the same procedure as for the inclusive proton events, and found to be 0.31 ± 0.22 . This is the first derivation of the α_p^{NM} value for the exclusive process $\Lambda p \rightarrow np$.

The threshold-energy dependence of the α_p^{NM} values




FIG. 2: (a) Opening-angle distribution of a neutron and a proton from the non-mesonic decay of ${}^5_\Lambda\text{He}$. Back-to-back events are clearly seen. (b) An excitation energy spectrum with the proton-neutron pair emitted in the back-to-back direction.

TABLE II: Summary table.

		$2^\circ \sim 6^\circ$	$6^\circ \sim 9^\circ$	$9^\circ \sim 15^\circ$
(π^+, p)	A_p	-0.000 ± 0.002	0.003 ± 0.002	0.003 ± 0.002
	A_π	-0.001 ± 0.001	0.003 ± 0.001	0.000 ± 0.001
(π^+, K^+)	P_Λ	0.149 ± 0.069	0.266 ± 0.076	0.694 ± 0.076
	α_p^{NM}		$0.11 \pm 0.08 \pm 0.04$	
$E_p^{th} = 40 \text{ MeV}$	α_p^{NM}		$0.12 \pm 0.09 \pm 0.05$	
$E_p^{th} = 60 \text{ MeV}$			$0.20 \pm 0.10 \pm 0.05$	
$E_p^{th} = 80 \text{ MeV}$			$0.09 \pm 0.14 \pm 0.04$	
np coincidence	α_p^{NM}		0.31 ± 0.22	

was further scrutinized by changing the proton detection energy thresholds as 40, 60, and 80 MeV (see Table 2). When protons are coming from the $\Lambda p \rightarrow np$ process, the proton energy distribution should have a broad bump structure peaked at ~ 75 MeV. The effects of FSIs and two-nucleon induced decay mode ($\Lambda p N \rightarrow np N$) would contribute mainly at low energy and reduce the proton decay asymmetry. In a recent calculation by Alberico et al. [13], it was demonstrated that the α_p^{NM} central values were reduced by 16% with the 30 MeV threshold, as compared to that with the 70 MeV threshold, due to the FSI effect introduced in their calculation. On the contrary, we obtained rather stable α_p^{NM} values.

As for theoretical models based on the one-meson-exchange mechanism, the introduction of one-kaon-exchange increased the contribution of the amplitude f in Table I, and solved the Γ_n/Γ_p puzzle. However, at the same time it increased the contribution of the third term in Eq. 2, and came up with a large negative value in

α_p^{NM} of $-0.6 \sim -0.7$. In this framework, it seems to be very hard to resolve the situation. A recent calculation of K. Sasaki *et al.* [14] suggests that there could be a solution by further introducing the one-sigma-exchange mechanism of a scalar-isoscalar meson σ . It reduces the effect of the third term in Eq. 2 without changing the branching ratios very much.

In summary, we obtained the proton asymmetry parameter in the non-mesonic weak decay of the polarized ${}^5_\Lambda\text{He}$ hypernucleus in the exclusive process $\Lambda p \rightarrow np$. The process was selected by observing the proton-neutron pairs emitted in the back-to-back kinematics in coincidence. The obtained value of $\alpha_p^{NM}=0.31\pm0.22$, together with that for the inclusive protons, $\alpha_p^{NM}=0.11\pm0.08\pm0.04$, largely deviates from the theoretical values of around -0.6 , although these theoretical models well reproduce the Γ_n/Γ_p ratio. The stability of the experimental α_p^{NM} values with different proton energy thresholds was also confirmed. This seems to suggest the importance of a scalar-type meson-exchange mechanism, such as a scalar-isoscalar σ meson or a correlated π - π pair.

We would like to thank Prof. K. Nakamura and the KEK-PS staff for their support to the present experiment. We also acknowledge the staff members of the cryogenic group at KEK. This work was supported in part under the Korea-Japan collaborative research program of KOSEF(R01-2005-000-10050-0) and KRF(2003-070-C00015).

* Present address: Osaka Electro-Communication University, Osaka 572-8530, Japan

- [†] Present address: Korea Research Institute of Standards and Science (KRISS), Daejeon, 305-600, Korea
- [‡] Present address: RIKEN, Wako, Saitama 351-0198, Japan
- [§] Present address: Japan Atomic Energy Research Institute, Tokai, Ibaraki 319-1195, Japan
- [¶] Present address: High Energy Accelerator Research Organization (KEK), Tsukuba, Ibaraki 305-0801, Japan
- [1] M.M. Block and R.H. Dalitz, *Phys. Rev. Lett.* **11**, 96 (1963).
- [2] H. Nabetani, T. Ogaito, T. Sato and T. Kishimoto, *Phys. Rev. C* **60**, 017001 (1999).
- [3] W.M. Alberico, G. Garbarino, *Phys. Rep.* **369**, 1-109 (2002), and references therein.
- [4] H. Ota *et al.*, *Nucl. Phys.* **A754**, 157 (2005) ; B.H. Kang, Ph.D. thesis, Seoul National University, 2004.
- [5] A. Parreno, A. Ramos, and C. Bennhold, *Phys. Rev. C* **56**, 339 (1997).
- [6] K. Sasaki, T. Inoue, and M. Oka, *Nucl. Phys.* **A707**, 477 (2002).
- [7] S. Ajimura *et al.*, *Phys. Rev. Lett.* **84**, 4052 (2000).
- [8] T. Fukuda *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **361**, 485 (1995).
- [9] H. Bando, T. Motoba, M. Sotona and J. Zofka, *Phys. Rev. C* **39**, 587 (1989).
- [10] S. Okada *et al.*, *Phys. Lett. B* **597**, 249 (2004).
- [11] T. Motoba and K. Itonaga, *Nucl. Phys.* **A577**, 293c (1994).
- [12] S. Ajimura *et al.*, *Phys. Rev. Lett.* **80**, 3471 (1998).
- [13] W.M. Alberico, G. Garbarino, A. Parreno and A. Ramos, *Phys. Rev. Lett.* **94**, 082501 (2005).
- [14] K. Sasaki, M. Izaki, and M. Oka, *Phys. Rev. C* **71**, 035502 (2005).